


Chapter 17 Food Protection with High-Temperature and Characteristics of Thermophilic Microorganisms

Tsuei-Yun Fang

節錄自 Modern Food Microbiology, Jay, J. M., 7th ed.



Introduction

- ◆ Based on the destructive effect of high-temperature on microorganisms.
- ◆ **Pasteurization** → either the destruction of **all disease-producing organisms** (for example, **pasteurization of milk**) or the **destruction or reduction in the number of spoilage organisms** in certain foods, as in the **pasteurization of vinegar**.



Introduction

- ◆ The **pasteurization of milk** is achieved by heating as follows:
 - **145°F (63°C) for 30 min (low temperature long time, LTLT)**
 - **161°F (72°C) for 15 sec (high temperature short time, HTST, method)**
 - **191°F (89°C) for 1.0 sec**
 - 194°F (90°C) for 0.5 sec
 - 201°F (94°C) for 0.1 sec
 - 212°F (100°C) for 0.01 sec
- ◆ **These treatments are equivalent and are sufficient to destroy the most heat resistant of the nonsporeforming pathogenic organisms**



Introduction

- ◆ Milk pasteurization temperatures are sufficient to destroy **all yeasts, molds, Gram-negative bacteria, and many Gram-positives.**
- ◆ Two groups of organisms that survive milk pasteurization are: **thermodurics (耐熱生物)** and **thermophiles (嗜熱生物).**



Introduction

- ◆ **Thermoduric organisms** are those that **can survive** exposure to relatively high temperatures but **do not necessarily grow** at these temperatures.
- ◆ The nonsporeforming organisms that survive milk pasteurization generally belong to the genera *Streptococcus* (鏈球菌屬) and *Lactobacillus* (乳桿菌屬).



Introduction

- ◆ **Thermophilic organisms** are those that **not only survive** relatively high temperatures but **require high temperatures** for their growth and metabolic activities.
- ◆ The genera *Bacillus* (芽孢桿菌屬) and *Clostridium* (梭狀桿菌屬) contain the thermophiles of greatest importance in foods.



Introduction

- ◆ **Sterilization: the destruction of all viable organisms**
- ◆ Canned foods are sometimes called "**commercially sterile**" to indicate that **no viable organisms can be detected** by the usual cultural methods employed or that **the number of survivors is so low** as to be of no significance under the conditions of canning and storage.



Introduction

- ◆ The processing of **milk and milk products** can be achieved by the use of **ultrahigh temperatures** (UHT).
- ◆ The primary features of the UHT treatment include **its continuous nature** → requiring **aseptic storage and aseptic handling** of the product downstream from the sterilizer
- ◆ The **very high temperatures** (in the range **140-150°C**) and the correspondingly **short time (a few seconds)** necessary to achieve commercial sterility.



Introduction

- ◆ **UHT-processed milks have higher consumer acceptability** than the conventionally heated pasteurized products, and because they are **commercially sterile**, they may be stored at room temperatures for up to 8 weeks without flavor changes.

「超高溫瞬間殺菌法」(UHT-pasteurization, 120~130 °C, 2~5 秒), 國內市售鮮乳約 90% 均使用 UHT 設備進行殺菌程序, 僅少數乳品加工廠採用 LTLT、HTST 或合併膜過濾的殺菌方式。依現行法令規定, 超高溫瞬間殺菌乳 (UHT milk) 如採全程冷藏保存, 即可標示為鮮乳; 如經滅菌處理及無菌包裝, 可於常溫保存, 則須標示為保久乳 (農委會)



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

1. Water.


- The **heat resistance** of microbial cells **increases with decreasing humidity or moisture** (Table 17-1).
- **Protein denaturation** may be involved in the **mechanism of death by heat**
 - Heating of wet proteins → formation of free —SH groups → water-binding capacity of proteins increased → thermal breaking of peptide bonds easier



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

2. Fat (triglyceride, triesters of glycerol)

- Increase in the heat resistance of some microorganisms (Table 17-2). → **fat protection** ← directly affecting cell moisture.
- Heat-protective effect of **long-chain fatty acids is better** than that of short-chain acids.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

3. Salts.

- Depend on the kind of salt and concentration employed
- **NaCl** could **decrease water activity** and thereby **increase heat resistance** by a mechanism similar to that of drying, whereas **others may increase water activity** (e.g. Ca^{2+} and Mg^{2+}) and, consequently, **increase sensitivity to heat**.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

4. Carbohydrates.

- Sugars causes an **increase in the heat resistance** of microorganisms → may due to the **decrease in water activity** caused by high concentrations of sugars.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

5. pH.

- Microorganisms are **most resistant to heat at their optimum pH of growth**, which is generally about 7.0. As the pH is lowered or raised from this optimum value, there is a consequent increase in heat sensitivity (Fig. 17-1).
- **High-acid foods require less heat** to achieve sterilization compared to foods at or near neutrality.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

6. Proteins and Other Substances.

- Proteins have a **protective effect** on microorganisms. → high-protein-content foods must be heat processed to a greater degree than low-protein-content foods
- The presence of **colloidal-sized particles** in the heating foods also offers protection against heat.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

7. Numbers of Organisms.

- The **larger the number** of organisms, the **higher is the degree of heat resistance** (Table 17-3).
- Heat protection by large microbial populations → may be due to the **production of protective substances**, probably protein, excreted by the cells.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

8. Age of Organisms.

- **Most resistant to heat → stationary phase of growth (old cells)**
- Less resistant → logarithmic phase.
- **Heat resistance is also high** at the beginning of the **lag phase**
- **Old bacterial spores are more heat resistant** than young spores.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

9. Growth Temperature.

- The **heat resistance of microorganisms increase as the growth temperature increases**, and this is especially true for sporeformers.
- *Salmonella senftenberg* grown at 44°C was found to be approximately three times more resistant than cultures grown at 35°C (Table 17-4).



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

10. Inhibitory Compounds.

- A decrease in heat resistance ← presence of heat-resistant antibiotics, SO₂, and other microbial inhibitors.
- **Heat plus antibiotics and heat plus nitrite** → more effective in controlling the spoilage
- Adding inhibitors to foods prior to heat treatment → reduce the amount of heat.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

11. Time and Temperature.

- Higher the temperature → greater the killing effect of heat
- As **temperature increases** → **heating time decreases**.
- The **size of the heating vessel or container** and its **composition** (glass, metal, plastic) is also important. It takes longer to effect pasteurization or sterilization in large containers than in smaller ones.



FACTORS AFFECTING HEAT RESISTANCE IN MICROORGANISMS

12. Effect of Ultrasonics.

- The exposure of bacterial endospores to ultrasonic treatments just before or during heating → **lower spore heat resistance.**



RELATIVE HEAT RESISTANCE OF MICROORGANISMS

- ♦ **Related to their optimum growth temperatures.** Psychrophilic microorganisms are the most heat sensitive, followed by mesophiles and thermophiles.
- ♦ **Sporeforming bacteria are more heat resistant** than nonsporeformers, and **thermophilic sporeformers are more heat resistant** than mesophilic sporeformers.



RELATIVE HEAT RESISTANCE OF MICROORGANISMS

- ◆ **Gram-positive bacteria are more heat resistant** than gram-negative.
- ◆ The **asexual spores of molds** are slightly more heat resistant than mold mycelia.
- ◆ **The extreme heat resistance of bacterial endospores** is of great concern in the thermal preservation of foods.



RELATIVE HEAT RESISTANCE OF MICROORGANISMS

- ◆ Endospore resistance is believed to be due to three factors: **protoplast dehydration**, **mineralization** (礦化作用: 有機物轉換成無機物), and **thermal adaptation**. Protoplast dehydration appears to be the primary factor.



THERMAL DESTRUCTION OF MICROORGANISMS

- ◆ **Thermal Death Time (TDT):** This is the **time necessary to kill a given number of organisms at a specified temperature**. By this method, the temperature is kept constant and **the time necessary to kill all cells is determined**.



THERMAL DESTRUCTION OF MICROORGANISMS


General procedure for determining TDT:

1. Place a **known number of cells or spores** in a sufficient number of sealed containers.
2. The organisms are then **placed in an oil bath and heated for the required time period**.
3. At the end of the heating period, containers are **removed and cooled quickly in cold water**.
4. The organisms are then **placed on a suitable growth medium**.
5. **Grow and then plate count**. Death is defined as the inability of the organisms to form a visible colony.

THERMAL DESTRUCTION OF MICROORGANISMS

- ◆ **Thermal Death Point:** the temperature necessary to kill a given number of microorganisms in a fixed time, usually 10 min.
- ◆ **D Value:** the decimal reduction time (九成滅菌時間), or the time required to destroy 90% of the organisms.
 - = minutes required for the destruction curve to traverse one log cycle
 - = $1/(\text{the slope of the destruction curve})$ (Fig. 17-2).
 - When D is determined at 250°F, it is often expressed as Dr.

$$\text{slope} = 1/D \rightarrow D = 1/\text{slope}$$






THERMAL DESTRUCTION OF MICROORGANISMS

- ◆ D value is the measure of death rate of microorganisms
- ◆ **D value reflects the resistance of an organism to a specific temperature** and can be used to **compare the relative heat resistance** among different organisms/spores
- ◆ D value for the same organism varies depending on the food type
- ◆ D value **is lower in acid foods** and **higher in presence of high proteins**



THERMAL DESTRUCTION OF MICROORGANISMS

- ◆ **Thermal Death Time Curve.**
 - A curve used to calculate z value (Fig. 17-3).
 - **D value in minutes** is plotted on the semilog paper along the log scale (y-axis), and **the temperature (in Fahrenheit) of heating** is plotted along the linear axis (x-axis).



THERMAL DESTRUCTION OF MICROORGANISMS

z Value:

- ◆ **the number of degree temperature (in Fahrenheit)** required for the thermal death time curve to pass through one log cycle.
= $1 / (\text{slope of the TDT curve})$ (Fig. 17-3).
- z provides information on the **relative resistance of an organism to different destructive temperatures.**



THERMAL DESTRUCTION OF MICROORGANISMS

◆ z Value:

- ◆ Example: If adequate heat process is achieved at 150°F for 3 min and Z value was determined as 17°F, the equivalent heat process will be either 30 min at 133°F or 0.3 min at 167°F.

F Value.

- ◆ This value is **the equivalent time (in min)** at 250°F (121°C) of all heat considered → to **destroy a specific number of spores or vegetative cells** of a particular organism.
- ◆ $F_0 = D_r(\log a - \log b)$, where a, b are the numbers of cells in the initial and final populations, respectively.



12-D concept

- ◆ 12D concept is used mainly **in low acid canned foods (pH >4.6)** where ***Clostridium botulinum*** is a serious concern
- ◆ The minimum heat process that reduce the probability of survival of the most resistant ***C. botulinum* spores to 10⁻¹² in canned food.**



SOME CHARACTERISTICS OF THERMOPHILES

- ◆ The growth temperature of **thermophiles** has a minimum of around 45°C, an optimum between 50°C and 60°C, and a maximum of 70°C or above.
- ◆ Five important thermophile genera in foods: *Bacillus* (芽孢桿菌屬), *Alicyclobacillus* (環脂酸芽孢桿菌屬), *Geobacillus* (土芽孢桿菌屬), *Clostridium* (梭菌屬, 又稱梭狀芽孢桿菌屬), and *Thermoanaerobacterium* (熱厭氧桿菌屬).



SOME CHARACTERISTICS OF THERMOPHILES

1. Enzymes

- ◆ Three groups of thermophilic enzymes:
 - Enzymes are **stable at the temperature of production** but can be inactivated at higher temperatures (e.g., ATPase).
 - Enzymes are **inactivated at the temperature of production in the absence of specific substrates** (e.g., catalase).
 - Enzymes are **highly heat resistant** (e.g., alpha-amylase).



SOME CHARACTERISTICS OF THERMOPHILES

1. Enzymes

- ◆ **Enzymes obtained from thermophiles are more heat resistant** than those enzymes from mesophiles.
- ◆ Possible reasons for enzymes become heat resistant: **higher levels of hydrophobic amino acids** and **binding of metal ions**, such as divalent ions (Mg^{2+}).



SOME CHARACTERISTICS OF THERMOPHILES

2. Ribosomes



- ◆ Thermal stability of ribosomes is correlated to the maximum growth temperature of microorganisms.
- ◆ The **increased G-C content in RNA** → **more stable structure** through more extensive hydrogen bonding → higher thermal resistance.



SOME CHARACTERISTICS OF THERMOPHILES

3. Flagella

- ◆ **Thermophilic flagella are more heat stable** than those of mesophiles, for example, thermophilic flagella stay intact at temperatures as high as 70°C; the mesophilic flagella fall off at 50°C.
- ◆ **Thermophilic flagella are more resistant to urea and acetamide** → suggesting more **effective hydrogen bonding** occurs in thermophilic flagella .



OTHER CHARACTERISTICS OF THERMOPHILIC MICROORGANISMS

1. **Nutrient Requirements: Thermophiles have a higher nutrient requirement** than mesophiles when growing at thermophilic temperatures.
2. **Oxygen Tension:**
 - As the **growth rate of microorganisms increases** → the **oxygen demand increases**. But the decreased solubility of oxygen cannot meet the demand.
 - This is one of the most important limiting factors of thermophilic growth in culture media.



OTHER CHARACTERISTICS OF THERMOPHILIC MICROORGANISMS

3. Cellular Lipids.

- The state of cellular lipids affects thermophilic growth.
- Mesophiles growing above their maximum range showed **decreases in lipid content** and **more lipid saturation**. → **saturated branched-chain fatty acids** are preferred
- The **decrease in the proportion of unsaturated fatty acids** ← **growth temperatures increase**



OTHER CHARACTERISTICS OF THERMOPHILIC MICROORGANISMS

4. Cellular Membranes.

- ♦ The **leakage** of ultraviolet light-absorbing and other material from cells undergoing "**cold shock**" → membrane related to high-temperature death.
- ♦ Most animal die when body temperatures reach between 40°C and 45°C and most psychrophilic bacteria are killed at about this temperature range → may due to the **melting of lipid constituents** of the cell or cell membrane.
- ♦ **Cellular membrane integrity** → critical to growth and survival at thermophilic temperature



OTHER CHARACTERISTICS OF THERMOPHILIC MICROORGANISMS

5. Effect of Temperature.

- ◆ Thermophiles do not grow as fast at their optimum temperatures as mesophiles do.
- ◆ **Thermophile enzymes are inherently less efficient** than mesophiles because of thermal stability ← discard growth efficiency to survive

6. Genetics.

- ◆ The genetic loci for streptomycin resistance and that for growth at 55°C were closely linked.
- ◆ The precise mechanisms about the high-temperature phenomenon remain mystery.



CANNED FOOD SPOILAGE

- ◆ Due to the following improper handling: **underprocessing, inadequate cooling, leakage through seams, and preprocess spoilage.**
- ◆ The spoilage of canned food can be **classified** based on acidity.



CANNED FOOD SPOILAGE

1. Low acid food

a) $4.6 < \text{pH} < 6.8$



b) Found in meat and marine products, milk, some vegetables (corn, lima beans), meat and vegetable mixtures, and so on.

c) Spoiled by

- **thermophilic flat-sour group:** *Geobacillus stearothermophilus*, *Bacillus coagulans*
- **sulfide spoilers:** *Clostridium nigrificans*, *C. bifermentans*
- **gaseous spoilers:** *Thermoanaerobacterium thermosaccharolyticum*



CANNED FOOD SPOILAGE

2. Acid food

a) $3.7 - 4.0 < \text{pH} < 4.6$



b) Found in fruits such as tomatoes, pears, and figs.

c) Spoiled by




- **Thermophilic spoilers:** *B. coagulans* types
- **Mesophiles:** *B. polymyxa*, *Paenibacillus macerans*, *C. pasteurianum*, *C. butyricum*, lactobacilli, and others.



CANNED FOOD SPOILAGE

3. High acid food



- a) **pH < 4.0-3.7**
- b) Found in fruits, and fruit/vegetable products, such as grapefruit, pickles, and so forth.
- c) Spoiled by **nonsporeforming mesophiles: yeasts, molds, lactic acid bacteria**



CANNED FOOD SPOILAGE

Canned food spoilage organisms may be characterized as: (also see Table 17-15)

1. Mesophilic organisms

- Putrefactive (致腐敗的) anaerobes
- Butyric anaerobes
- Aciduric flat sours
- Lactobacilli
- Yeasts
- Molds

2. Thermophilic organisms

- Thermophilic anaerobes producing sulfide
- Flat-four spores
- Thermophilic anaerobes not producing sulfide



Appearance of canned food spoilage

- ◆ The normal appearance is **flat** or **slightly concave** (凹).
- 1. **Flipper** (急跳罐，輕度膨罐):
 - the can looks normal but one end of the can be made convex (凸) by striking or heating the can.
 - Not always represent microbial spoilage.
- 2. **Springer** (彈性罐):
 - both ends may be bulged but one can be pushed back into place, or else a can with one end bulged that when pressed in, pops out the other end.
 - Not always represent microbial spoilage.



Appearance of canned food spoilage

- 3. **Soft swell** (軟膨罐):
 - Both ends bulged that may be dented by pressing with the fingers.
 - Often represent **microbial spoilage**.
- 4. **Hard swell** (硬膨罐):
 - Both ends bulged, so that neither end can be dented by hand.
 - Often represent **microbial spoilage**.
 - In high-acid foods, it is referred to **hydrogen swells** (氫脹罐), which result from the release of hydrogen gas.
 - Two other common gases in cans of spoiled foods are **CO₂** and **H₂S**



Appearance of canned food spoilage

5. "Leakage-type" spoilage:

- The organisms enter cans **at the start of cooling through faulty seams**, which generally result from **can abuse**.
- The organisms that cause leakage-type spoilage can be found either on the cans or in the cooling water.
- This problem is **minimized** if the cannery **cooling water contains <100 bacteria/ml**.